



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

KALA AZAR AND BED BUGS

There has been a disposition for some years to seek to fasten on the bed-bug the responsibility for transmitting some of the infectious diseases. The habits of the animal lend decided substance to these expectations and the odium in which it is held doubtless makes its case the more hopeless.

Capt. W. S. Patton, I. M. S., (Ind. Med. Gaz., Feb. 19, 1912), reports that he has, by feeding *Cimex rotundatus* and *C. lectularius* on a case of Kala Azar, in whose blood the parasite had been identified, discovered an extended development of the parasite in these bugs. He also discovered that in bugs, in which the developing forms are present, again fed on the blood of a Kala Azar case, the flagellates are destroyed within 24 hours. However, if the bugs are not fed the second time the development goes on and is completed by the 10-12th day after the single feed. The parasite goes thru its complete cycle in both species. By the 7th-9th day the bug swarms with the flagellate stage in masses of rosettes. It is also claimed that the bugs do not contain, naturally, a flagellate.

CELL SIZE AND NUCLEAR SIZE

Conklin (Jour. Exp. Zool. Jan. 1912) in an elaborate paper based on a series of experiments and observations on *Crepidula* and *Fulgar* gives a most interesting series of conclusions concerning the size of cells and nuclei. Among these may be mentioned:

1. Inequality of cell division is due to internal causes rather than to pressure from without; and the controlling factors within are more intimate than the mere presence of metabolic substances such as yolk.
2. The size ratio between plasma and nucleus in the different blastomeres of an egg is not constant nor self-regulating; but seems to depend rather on the rate of cell division.
3. The inciting cause of cell division seems to lie in the coincidence of centrosomal, chromosomal, and cytosomal rhythms.
4. The size of the nucleus is dependent upon at least three factors: the initial quantity of chromatin; the volume of the cytoplasm; the length of the resting period.
5. In *Crepidula* the volume of cytoplasm more than doubles during the passage from the 1-cell stage to the 24-cell stage; the

yolk decreases in volume nearly one-half in that time; and the entire embryo has less mass at the 24 than at the 1-cell stage. This shows that the protoplasmic growth during cleavage occurs at the expense of the yolk.

6. The average increase of protoplasm for each division in early cleavage (32 cells) is 6 per cent; of nucleus, 5-9 per cent; of chromatin, 8 per cent.

PROTOZOA IN HAY INFUSIONS

Woodruff (Jour. Exp. Zool., Feb., 1912), discusses a number of series of experiments on hay infusions with a view to determining the sources and sequence of their protozoan population. To determine the sources, he (1) used sterilized hay and water, exposed to air; (2) sterilized water with fresh hay, and air excluded; (3) ordinary tap water, with sterilized hay and air excluded; and (4), as controls, fresh hay, tap water, and loosely covered vessels. The experiments, which are valuable to the ordinary laboratory worker, furnish the following conclusions:

1. Air, water, and hay are all sources of protozoa in infusions,—the air being least, and the hay most, important. But ordinary hay, added to ordinary tap water, while furnishing some protozoa, will not produce a sufficient number of representative protozoa for the study of the full sequences.

2. In order, therefore, to study standard sequences it is necessary to "seed" the infusions with matter from general laboratory cultures. In such "seeded" infusions a definite sequence of appearance, of dominance (or maximum) and of disappearance was observed. The sequence of appearance at the surface of the infusion is as follows: Monad, Colpoda, Hypotrichida, Paramecium, Vorticella, and Ameba.

3. The middle of the infusion is inhabited chiefly by free-swimming types brought there by over crowding at the top and bottom, and does not manifest so definite a sequence.

4. The appearance, in appreciable numbers, of any of these types (except Ameba), at the bottom of the infusion, coincides with, or quickly follows, its surface maximum,—and seems to indicate the beginning of its decline.

5. As biological elements entering into the determination of